



Association of Minimal Access Surgeons of India \*



Indian Association of Gastrointestinal Endo Surgeons \*



Society of Endoscopic and Laparoscopic Surgeons of India

**INTER ASSOCIATION SURGICAL PRACTICE RECOMMENDATIONS IN COVID 19 ERA**  
**(FOR MINIMAL ACCESS SURGEONS IN INDIA)**

23<sup>rd</sup> APRIL 2020

In the current pandemic COVID 19 scenario, there is urgent need for guidelines for performing various surgical procedures. The utmost priority remains the safety of the patients and health care workers. Current recommendations are mostly based on expert opinion and knowledge of other pathogens with similar characteristics. These recommendations are time sensitive. These are bound to change as evidence-based data emerge.


General recommendations:

- We should provide timely surgical care to patients presenting with urgent and emergent surgical conditions while optimizing patient care resources (e.g. hospital and intensive care unit beds, personal protective equipment, ventilators) and preserving the health of caregivers.
- All elective surgical and endoscopic cases should be postponed at the current time.
- Non-operative management should be considered where ever possible (such as for early appendicitis and acute cholecystitis).
- All non-urgent in-person clinic/office visits should be cancelled or postponed, unless needed to triage active symptoms or manage wound care.
- All non-essential hospital or office staff should be allowed to stay home and telework.
- Multidisciplinary team (MDT) meetings should be sought after in any surgical decision and should be held virtually as possible. This team include the surgeon, intensivist, radiologist, infectious disease specialist and nurse managers. (Subject to availability and requirements)
- In case the treating hospital is not a COVID 19 designated center, patients with high risk and diagnosed cases should be referred to appropriate designated COVID 19 center.

Surgical Consultation recommendation:

- Avoid physical consultation for non-emergency cases.
- Use the telemedicine platform or provider's help and consult your patients online for normal surgical problems and follow-up. Please follow the regulations issued by the MOHFW. <https://www.mohfw.gov.in/pdf/Telemedicine.pdf>
- Please follow the statutory policies of respiratory triaging in receiving surgical emergencies.
- The social distancing, restricting the attendants to not more than one, patient/attendant masking and hand hygiene measures should be adhered to.
- The entry and exit pathway should be according to statutory regulations.
- It is desirable, all preoperative patients be tested for COVID 19 status a day prior to surgery.
- In referral or transfer from other health care facility - Please discuss with the referring Physician in detail about the contact details, overseas travel history of the patient or family members and presence of respiratory symptoms.

**Surgical Triage:**



**Definitions:**

- **Emergency Life threatening conditions-** these are those conditions in which immediate surgical intervention is required and there is not enough time available to do and get the results of COVID 19 testing. E.g. – life threatening Traumatic hemoperitoneum with hemodynamic instability, bowel gangrene.
- **Emergency Procedures-** These are those procedures that require immediate surgical intervention but provide a window of few hours to get COVID 19 testing. Interim non-surgical intervention could also be an option. E.g.- Bowel obstruction, perforation of hollow viscus. Emergency procedures are undertaken in life threatening conditions and have no alternatives, e.g. bowel perforation, gangrene and unresolved obstruction, complicated appendicitis, complicated cholecystitis, obstructed / strangulated hernia which need immediate surgery.

- Semi emergent procedures- These includes condition where the procedure can be deferred for a few weeks or months but not more than 3 months due to worsening of symptoms or progression of stage of disease affecting final outcome. E.g.- Major malignancies.
- Elective Procedures- These include those procedures which can be postponed by 3 months or more with mutual consent between patient and surgeon with no untoward effect on final outcome. E.g.- Uncomplicated groin hernia

#### Pre operative Phase:

- All patients in addition to surgical evaluation should be tested for COVID 19 RT PCR where possible.
- In emergency situations, patient should have a Xray Chest. CT Chest is done if there is a need for CT abdomen.
- Non-surgical percutaneous interventions are prudent if it can defer surgery for the time being and minimize hospital stay and requirement for a longer institutionalized health care.
- Where possible, the role of alternate non-surgical approaches to GI cancer such as neoadjuvant therapy and radiation therapy are to be considered. A virtual multidisciplinary approach is advised.



#### Consent Directives:

- During COVID 19, an improvised consent form with additional information about COVID 19 and its associated risks have to be obtained.
- A model consent form is attached as an annexure.

#### Personal Protection Equipment:

- The personnel in OT should don adequate Personal Protective Equipment. PPE GUIDELINES From the MOHFW are available at:  
<https://www.mohfw.gov.in/pdf/GuidelinesonrationaluseofPersonalProtectiveEquipment.pdf>


- Donning of PPE shall be done in procedure room/ OT Room and Doffing shall be done in the designated wash area.



Operating Room Directives:

1. Negative Pressure room is desirable where possible
2. Minimum no of personnel to be inside operating room
3. Limit the size of surgical team as much as possible
4. Avoid the operating room personnel stepping out of OT during the procedure
5. Minimum 1-hour time gap to be given between two procedures / surgeries.
6. Minimize duration of surgery
  - a. No surgical or nursing training during this period
  - b. Avoid multiple and complex procedures
7. Reusable accessories are cleaned and disinfected with appropriate solutions as soon as the procedure is over.
8. 1% Sodium Hypochlorite solution cleaning is recommended for OT Tables and trolleys as soon as the patient is shifted.

Flow Chart COVID Infection Prevention & Control Policy in Operating Room



### Anesthesiologists Directives:

The regulations for the anesthesiologists are issued by the Indian Society of Anesthesiologists. Salient outcomes are as follows:

- Regional anesthesia is preferred.
- GA with ET intubation is high aerosol producing event.
- The surgical team is advised to enter OT minimum fifteen minutes after the patient has been anesthetized
- Laminar flow / Air conditioners at the OT is to be started after induction of anesthesia.
- Stop laminar airflow / air conditioners twenty minutes before extubation.
- Post-surgery all anesthesia equipment to be cleaned with 1% sodium hypochlorite solution.

### Intra operative recommendations for Laparoscopic Surgery

Pneumoperitoneum and Aerosol Protection:

Currently, the best practice for mitigating possible infectious transmission during open, laparoscopic and endoscopic intervention is to use a multi-faceted approach, which includes:

1. Proper room filtration and ventilation
  2. Appropriate PPE
  3. Smoke evacuation devices with a suction and filtration system
- Once placed, ports should not be used to evacuate smoke or de-sufflation during the procedure without adequate precautions.
  - If movement of the insufflating port is required, the port should be closed prior to disconnecting the tubing and the new port should be closed until the insufflator tubing is connected. The insufflator should be "on" before the new port valve is opened to prevent gas from back-flowing into the insufflator.

- During desufflation, all escaping CO<sub>2</sub> gas and smoke should be captured with an ultra-filtration system and desufflation mode should be used on your insufflator if available.
- If the insufflator being used does not have a desufflation feature, be sure to close the valve on the working port that is being used for insufflation before the flow of CO<sub>2</sub> on the insufflator is turned off. Without taking this precaution contaminated intra-abdominal CO<sub>2</sub> can be pushed into the insufflator when the intraabdominal pressure is higher than the pressure within the insufflator.
- The patient should be flat and the least dependent port should be utilized for desufflation.
- Specimens should be removed once all the CO<sub>2</sub> gas and smoke are evacuated.
- Surgical drains should be utilized only if absolutely necessary.
- Suture closure devices that allow for leakage of insufflation should be avoided.
- The fascia should be closed after desufflation.
- Hand-assisted surgery should be avoided as it can lead to significant leakage.
- Wound protection devices used for specimen retrieval should only be placed after desufflation.
- The leakage around cannulas need to be minimised.
- Suction devices are a potential source of infection. Use of appropriate filter is advised.
- For routine evacuation of smoke and fumes, a member of the operating team should be designated for this and he/she should use the side channel for controlled evacuation. Standardised smoke evacuation devices can be used.



- All pneumoperitoneum should be safely evacuated from the port attached to the filtration device before closure, trocar removal, specimen extraction or conversion to open.

### Energy Devices:

“Aerosolization” of viral and bacterial RNA/DNA may occur during the use of energy devices in general surgery - both open and laparoscopic - although there is limited evidence that viable infective particles are dispersed.

The various energy sources lead to varying particle sizes with electrocautery and laser having the smallest, hottest particles and ultrasonic devices larger, cooler particles.

During both open and laparoscopic surgery, the particle concentration tends to increase over time of use of energy sources.

- Use of energy devices is a potential risk.
- Minimal use and short burst usage are advised.
- Judicious use of energy devices and smoke evacuation is advised.
- Cold hemostasis is the choice method advised.
- The energy sources should be used at the lowest power setting and charring of tissues should be avoided to minimize the creation of smoke.

### Flexible Endoscopy:

Surgical colleagues perform various flexible endoscopic procedures. It is pertinent to follow safe endoscopy practice directives issued by the respective society.

- We should stop all elective endoscopy work. Flexible endoscopy and therapeutic procedures are indicated only for clearly defined emergency and urgent cases.
- All Staffs in the unit should wear N 95 mask and good quality PPE during any endoscopic procedure
- Every emergency or urgent endoscopy patient should be presumed as Covid positive and managed accordingly taking all strict safety precautions

- Comprehensive well written informed consent form comprising of all necessary details relevant to covid pandemic should be signed by the patient and relative prior to undergoing any endoscopic procedure
- All emergency and urgent endoscopic procedures should preferably be done under GA with careful endotracheal intubation.
- Endoscopy requires high level disinfection while endoscopic accessories should be either disposable or need sterilization.

Conclusion:

Our surgical health care system should adapt to this rapidly changing health environment due to covid 19 pandemic. Next few months are going to be critical in mitigating the spread of infection. Surgical interventions are considered only for clearly defined emergency and urgent cases. We should understand our vital role in protecting our staffs while treating our surgical patients. We should ensure best surgical care by having clear management strategy and compassionate approach.



Prof Bhupinder Singh Pathania

President – AMASI



Dr Ramen G Goel

President – IAGES



Dr Virinder Kumar Bansal

President – SELSI



Dr Kalpesh V Jani

Secretary – AMASI



Dr S Easwaramoorthy

Secretary – IAGES



Dr Amit Srivatsava

Secretary – SELSI

Drafted and Reviewed By:

Dr Tamonas Chaudhuri – AMASI    Dr S Easwaramoorthy – IAGES    Prof Pawanindralal – SELSI

Dr Kalpesh Jani – AMASI    Dr M Kanagavel – IAGES    Dr Krishna Asuri – SELSI

DISCLAIMER: This document is not valid for Medico-legal purpose.

## References:

1. <https://www.sages.org/notes-from-the-battlefield-april-6-2020/>
2. <https://www.sages.org/notes-from-the-battlefield-march-30-2020/>
3. <https://www.sages.org/wp-content/uploads/2020/04/Closing-the-Back-Door-SOP.pdf>
4. <https://www.sages.org/recommendations-surgical-response-covid-19/>
5. Rothan H, Siddappa B. The epidemiology and pathogenesis of coronavirus disease (COVID-19) Outbreak. Journal of Autoimmunity. <https://doi.org/10.1016/j.jaut.2020.102433>. 2020.
6. Zheng M, Boni L, Fingerhut A. Minimally invasive surgery and the novel coronavirus outbreak: lessons learned in China and Italy, Annals of Surgery. 2020.
7. <https://www.sages.org/resources-smoke-gas-evacuation-during-open-laparoscopic-endoscopic-procedures/>
8. <https://journals.lww.com/annalsofsurgery/Documents/Are%20we%20harming%20cancer%20patients.pdf>
9. [https://www.isde.net/resources/Documents/Resources/ISDE\\_Position\\_Statement\\_COVID19\\_2020.03.30.pdf](https://www.isde.net/resources/Documents/Resources/ISDE_Position_Statement_COVID19_2020.03.30.pdf)
10. [https://umbraco.surgeons.org/media/5162/guidelines-for-emergency-upper-gi-and-bariatric-surgery-during-the-covid- v2\\_6-april.pdf](https://umbraco.surgeons.org/media/5162/guidelines-for-emergency-upper-gi-and-bariatric-surgery-during-the-covid- v2_6-april.pdf)
11. [https://umbraco.surgeons.org/media/5136/optimal-surgical-approach-during-the-covid-19-pandemic\\_updated-version.pdf](https://umbraco.surgeons.org/media/5136/optimal-surgical-approach-during-the-covid-19-pandemic_updated-version.pdf)
12. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC7117791/pdf/FVvinObGyn-12-3.pdf>
13. <https://www.facs.org/covid-19/clinical-guidance/review-committee>
14. <https://www.facs.org/covid-19/clinical-guidance/statement-maintaining>
15. <https://www.facs.org/covid-19/clinical-guidance/surgeon-protection#ppe-use>
16. <https://pbs.twimg.com/media/EUjMyWrWAAIt7bh.jpg>
17. <https://pbs.twimg.com/media/EUjMyWqWAAyxXhN.jpg>

**REFERENCE CONSENT FORM FOR SURGERY DURING COVID 19 PANDEMIC**

- I/We \_\_\_\_\_ knowingly and willingly consent to have treatment / surgery / investigation completed during the COVID-19 pandemic.
- I/We have been informed that I am opting for treatment / surgery / investigation in the midst of community outbreak of COVID19 pandemic.
- I/We have been explained about the risks and side effects of COVID 19 infection and precautions being taken by the hospital to prevent transmission of infection in patients and staff.
- I/We understand and acknowledge that although all precautions as prescribed by the Government of India have been duly implemented by the hospital, as the disease is new and its kinetics, symptoms and treatment are still being studied, no assurances or guarantees can be offered regarding disease transmission to the patient or clinical outcome in case of disease transmission to the patient.
- I/We understand the implications and give our consent for the continuation of treatment / surgery / investigation.
- I have been tested for COVID 19 to found be negative and I have been told that I may become positive in the post operative period, if I am in the carrier stage of the disease.
- I have not been tested and I agree to continue with the surgical procedure. If found positive later, i would not attribute the same to the surgical procedure conducted and the surgeon and doctor concerned.
- I have been explained that there is evidence that surgery in infected but asymptomatic patients is associated with a more severe disease manifestation in the post-operative period and an increased mortality.

Patient's Name: \_\_\_\_\_ Signature \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Doctor's Name: \_\_\_\_\_ Signature \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Witness's Name: \_\_\_\_\_ Signature \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Contact No. of witness \_\_\_\_\_ Address \_\_\_\_\_

Attendant's Consent: If patient is unable to give consent, reason (Minor/unconscious/under sedation/ mental incapacity/ disoriented, other please specify)

Attendant's Name: \_\_\_\_\_ Relationship \_\_\_\_\_

Signature of the Attendant: \_\_\_\_\_ Date \_\_\_\_\_ Time \_\_\_\_\_

Statement of Interpreter (where applicable):

I have comprehensively explained the information above along with the discussion/explanations provided by the doctors, to the patient and /or his/her attendant and in a way which I believe he/she/they can understand and the patient and /or his/her attendant have informed me that they have understood the aforesaid information completely.

Interpreter's Name \_\_\_\_\_ Signature \_\_\_\_\_ Contact No \_\_\_\_\_

Date \_\_\_\_\_ Time \_\_\_\_\_